

# Chemoreceptors of Crustaceans: Similarities to Receptors for Neuroactive Substances in Internal Tissues

by William E. S. Carr,\* Barry W. Ache,\* and Richard A. Gleeson\*

A description is given of crustacean chemosensory systems and the neurophysiological procedures used to study them. Their response properties and tuning characteristics are discussed. A review is then provided of specific crustacean chemoreceptors that are stimulated selectively by either purine nucleotides, taurine, glutamate, or glycine, all of which have neuroactive properties in internal tissues.

Two distinctly different types of purinergic chemoreceptors occur on the antennules of the spiny lobster.  $P_1$ -like chemoreceptors have a potency sequence of  $AMP > ADP > ATP > \text{adenosine}$  and show a strict structural requirement for the ribose phosphate moiety.  $P_2$ -like chemoreceptors have a potency sequence of  $ATP > ADP > AMP$  or adenosine and show a broad sensitivity to nucleotide triphosphates with modifications in both the purine and ribose phosphate moieties. Sensilla containing the dendrites of chemosensory neurons also possess an ectonucleotidase(s) that inactivates excitatory nucleotides to yield adenosine which is subsequently internalized by a sensillar uptake system.

Narrowly tuned taurinergic chemoreceptors are present on both the antennules and legs of lobsters. Although taurine itself is the most effective stimulant, the taurine analogs hypotaurine and  $\beta$ -alanine are also very excitatory. Structure-activity studies indicate these chemoreceptors have marked similarities to taurine-sensitive systems in internal tissues of vertebrates. By contrast, comparative studies of glutamatergic chemoreceptors on the legs of lobsters indicate response spectra different from those of the glutamate receptors in lobster neuromuscular junctions and the three classes of excitatory amino acid receptors identified internally in vertebrates. Crustacean chemoreceptors for glycine, ecdysteroids, and pyridine are also described. The hypothesis that receptors for internal neuroactive agents may have originally evolved as external chemoreceptors of primitive aquatic organisms is discussed.

## Introduction

The chemical senses of many aquatic animals are extremely well developed. Waterborne chemicals influence facets of behavior as diverse as food finding (1,2), predator avoidance (3,4), mate recognition and mating (5,6), substrate selection and metamorphosis by larvae (7,8), homing by migratory species (9,10), and species recognition and social behavior (11,12). Chemoreception in aquatic animals has been reviewed recently (3,13-15).

Chemosensory studies have revealed that many of the external chemical agents stimulating behavioral and physiological responses in aquatic invertebrates are the very same substances that function internally in higher organisms as either transmitters or modulators. This paper will briefly describe crustacean chemosensory systems and the physiological procedures used to study them. A description will then be provided of specific crustacean chemoreceptors that are stimulated by sub-

stances known to have neuroactive properties in internal tissues.

## Chemosensory Systems in Decapod Crustaceans

Because of their large size and easily accessible chemosensory organs, certain decapod crustaceans such as the Florida spiny lobster (*Panulirus argus*), the California spiny lobster (*P. interruptus*), the American lobster (*Homarus americanus*), and certain crayfish species (e.g., *Orconectes limosus*) have proven to be popular animals for neurophysiological studies of the chemosensory process. The functional organization of the olfactory system in these crustaceans has been reviewed recently (1,16). The olfactory organ consists of a tuft of sensilla, termed aesthetasc sensilla, present on the lateral filament of the antennule (Fig. 1). In the Florida spiny lobster, *P. argus*, each antennule has about 2000 of these sensilla, with each sensillum being approximately 1000  $\mu\text{m}$  in length and 40  $\mu\text{m}$  in diameter. Each is innervated by the cilia-bearing dendrites of an

\*C. V. Whitney Laboratory and Department of Zoology, University of Florida, Route 1, Box 121, St. Augustine, FL 32086.




FIGURE 1. (A) Head of the spiny lobster showing the lateral branches of the antennules with tufts of chemosensory aesthetasc sensilla. (B) Schematic of aesthetasc sensillum depicting dendritic terminals housed within the cuticular sheath.

estimated 350 bipolar sensory neurons (17). Axons from these olfactory cells project via the antennular nerve to the midbrain where they make synaptic contacts in the glomerular neuropile of the paired olfactory lobes (18,19).

In addition to the aesthetasc sensilla of the antennules, a variety of other morphological types of chemosensory sensilla have been identified on the mouth parts, claws and walking legs of decapod crustaceans (20–23). Functionally these other chemoreceptors are often considered to be comparable to the taste system of vertebrates (1,24). The chemical sensitivities of receptors on the distal segments (dactyls) of the walking legs have been especially well studied (25–29).

## Procedures for Neurophysiological Analysis of Crustacean Chemoreceptors

The crustacean chemoreceptor cell is a primary sensory neuron with a spiking axon that projects directly to the brain. The responses of these cells to chemicals can be quantified by extracellularly recording the neural discharge of the axon when chemicals are applied to the dendrite. Such recordings are accomplished by mounting an excised chemosensory appendage in a chamber that allows the dendrite-bearing sensilla to be superfused with chemicals. One such arrangement for stimulating and recording from crustacean chemoreceptor cells is illustrated in Figure 2. Continuous perfusion with oxygenated saline maintains the viability of the preparation.

Chemical stimulation of the preparation in Figure 2 is achieved by injecting small volumes (i.e., 20 to 300  $\mu$ L) of solutions, as a bolus, into a carrier flow of water that continuously flushes over the sensilla. This procedure introduces the stimulus with no disruption of flow concurrent with the arrival of the stimulus at the preparation. Depending on the volume of stimulant injected, the preparation can be exposed to either a brief (e.g., 3 sec) pulse, or to a prolonged plateau of a constant concentration of the stimulant. Conductivity and fluorometric analyses show that this procedure delivers reproducible profiles of stimulant that approximate a step function on their leading edge.

Extracellular recording from receptor cells is achieved by splitting fascicles from the nerve and applying suction electrodes en passant to the exposed axons. Chemosensitive units are identified as such by their altered frequency of discharge to brief pulses of search stimuli and their lack of response to identically applied control carrier water. Responses from single neurons are obtained either by fortuitous separation of fascicles containing only a single active unit or by window (voltage and/or time) discrimination of single units from multi-unit recordings. Single-cell status of all records can be verified by an on-line template matching scheme that involves passing the real time analog signal through a delay register to retain the pretriggered aspect of the waveform and superimposing the entire waveform of each spike that triggers the window discriminator on the synchronized display of an analog storage oscilloscope. Evoked responses (spike trains) can be quantified by electrically counting the output of the window discriminator if the parameter of interest is the total num-




FIGURE 2. Diagram of a typical stimulus delivery and recording apparatus used to analyze crustacean chemoreceptor cells. An excised, perfused antennular preparation is illustrated; similar systems have been used to study dactyl chemoreceptors. Additional details are given in the text.

ber of spikes. Alternatively, a microprocessor-based interval analysis can be performed on the same output if it is necessary to obtain other, time-dependent parameters of the spike train.

Now it is also possible to record intracellularly from crustacean chemoreceptor cells (30). This technique is only beginning to be applied to the particular types of receptor cells discussed in this review, but the technology is now available to examine the biophysical basis of stimulus-response coupling in these cells.

## Response Properties of Crustacean Chemoreceptors

Most crustacean chemoreceptor cells have very low levels of spontaneous activity ( $\leq 1$  Hz). Chemical stimuli cause these cells to show a tonic increase in the frequency of discharge, although the response often becomes slightly more phasic at higher stimulus concentrations (Fig. 3). In nature, most chemoreceptors are only exposed intermittently to a chemical stimulus, and the actual pattern of the response is strongly influenced by the dynamics of the access of the stimulus to the receptor surface. The lobster antennule, for example, is flicked reflexively at a rate of 1–2 Hz. This behavior appears to regulate the access of the stimulus to the tight cluster of sensilla; the stimulus is sampled as a series of discontinuous pulses that onset within about 100 msec. The afferent discharge of the sensory cells reflects this pulsatile stimulus profile as a series of brief, phasic bursts (31).

In a chemosensory cell, different stimulatory com-

pounds usually evoke responses that have similar temporal patterns, although instances where the discharge pattern may change have been reported (32). Perhaps constancy in the temporal pattern of discharge with different stimulants is to be expected. The use of different temporal patterns to encode different stimulants would make it difficult to distinguish the pattern for a particular stimulant from patterning introduced by short-term fluctuations in stimulus concentration. The pulsatile nature of odorants in the aquatic environment has been described (14).

The rate of discharge of chemoreceptor cells is typically a simple function of the stimulus concentration (Fig. 3). The response of single cells usually saturates over 2 to 3 log units of concentration, although populations of cells reach saturation over broader ranges of concentration. The latter is presumably indicative of range fractionation which enables the overall population to be sensitive to a wide range of concentrations while still retaining maximal sensitivity to changes in concentration.

A striking feature of some crustacean chemoreceptors is their extremely low threshold. Antennular chemoreceptors in the spiny lobster give discrete responses to picomolar concentrations of some amino acids (33). Dactyl receptors in the American lobster respond to as little as 35 femtomolar ammonia (25).

## "Tuning" Characteristics of Crustacean Chemoreceptors

"Tuning" refers to the number and the heterogeneity of the different compounds that are stimulatory to a




FIGURE 3. Effect of stimulus concentration on the neural discharge of a chemoreceptor cell from the antennule of the spiny lobster, *P. argus*. Shown are the extracellularly recorded spike trains evoked by a 2-sec exposure to taurine at the molar concentrations indicated, and to seawater (ASW). The traces faithfully reproduce the temporal pattern of the train, but not the waveform of the spike, which has been shaped electrically for maximum clarity. Note that concentration has little effect on the temporal pattern of the response, primarily altering the frequency and number of spikes in the train.

chemoreceptor cell; tuning characteristics of crustacean chemoreceptors have been reviewed recently (34). In crustaceans it is known that different types of cells contribute to the peripheral chemosensory organization. Chemoreceptor cells with narrow response spectra are found on the dactyls (26,35) and antennules (36–38) of lobsters. These cells are narrowly tuned to particular amino acids and nucleotides found in food organisms. Specific examples of these types of cells will be described later. Such narrowly tuned cells are often called “specialists,” following terminology originally used by Schneider et al. (39) for moths. Other cells are less narrowly tuned, but are still limited in the breadth of their response. For example, the amino acid-selective units in the dactyls of the crayfish are limited in their response to a particular structural class of molecules, but do not appear to be selective within the class (40). Still other receptor cells are more broadly tuned and are sometimes called “generalists,” again by the terminology proposed by Schneider et al. (39). Broadly tuned receptor neurons respond to an array of structurally different substances that belong to different functional classes of molecules. In some broadly tuned neurons, sensitivity to different molecules is not nec-

essarily distributed randomly across the receptor population. When relatively large numbers of potentially stimulatory compounds have been tested on individual receptor cells in insects, for example, the cells tend to fall into “reaction groups.” Each group is composed of cells having part, or all, of their response spectra more similar to that of other cells in the same reaction group than to cells in other groups (41).

## Purinergic Chemoreceptors

### Background

Cyclic adenosine 3',5'-monophosphate has long been known to function both as a potent chemoattractant in cellular slime molds (42) and as a second messenger in many physiological processes (43,44). Regarding vertebrate organisms, adenosine and its noncyclic nucleotides are known to play major roles as transmitters or modulators of physiological activities in the nervous system, the vascular system, and in the smooth muscle of many internal organs (45–51). The purinergic receptors that mediate responses to adenosine (Ado) and adenine nucleotides have been classified by Burnstock (52) as  $P_1$  and  $P_2$  types. Among the distinguishing features of these two receptor types are the following:  $P_1$ -type purinoceptors have a potency sequence of  $\text{Ado} \geq \text{AMP} > \text{ADP} \geq \text{ATP}$ , whereas the  $P_2$ -type has a potency sequence of  $\text{ATP} \geq \text{ADP} > \text{AMP} \geq \text{Ado}$ . It will be shown below that marine crustaceans possess olfactory chemosensory cells with marked similarities to both the  $P_1$ - and the  $P_2$ -type purinoceptors found in internal tissues.

### $P_1$ -like Purinergic Chemoreceptors

The first evidence that  $P_1$ -like chemoreceptors are present in crustaceans came from the behavioral observation that adenosine 5'-monophosphate (AMP) was a potent attractant for the grass shrimp, *Palaemonetes pugio*. Bioassays of AMP analogs suggested that the behavioral response of the shrimp is mediated by receptors akin to  $P_1$ -type purinoceptors (53).

Documentation of the existence of purinergic chemoreceptors in crustaceans was obtained in physiological studies with the spiny lobster, *P. argus*, by employing the procedures for single-cell, extracellular recording as described above for ablated antennular filaments. Chemosensory purinoceptors with a potency sequence of  $\text{AMP} > \text{ADP} > \text{ATP} > \text{Ado}$  are present in the lobster antennule (Fig. 4) (37); this potency sequence, measured physiologically, is the same as that measured behaviorally in the shrimp. Studies of structure-activity relationships (SAR) of AMP and 16 analogs (Fig. 5) show that AMP is the most potent stimulant, and all changes in its structure result in a significant diminution in activity. However the greatest diminution accompanies changes in the ribose phosphate moiety rather than the adenine moiety. This is seen in Figure 5 by noting that eight of the nine least active analogs are those with modifications in the ribose phosphate




FIGURE 4. (A) Dose-response relationships for adenine nucleotides and adenosine as measured physiologically in chemosensory neurons of the spiny lobster. (B) (Inset) Behavioral results obtained with same substances in a shrimp. In both cases the potency sequence was AMP > ADP > ATP > adenosine. Data points are mean values ( $\pm$  SEM). From Derby et al. (37).

moiety. 8-bromo-AMP, the least active adenine-modified analog, differs from the others by having a *syn*- rather than an *anti*-conformation (54). Both the physiological response of the lobster chemoreceptors and the behavioral response of the shrimp are antagonized by the methylxanthine, theophylline.

Vertebrate  $P_1$ -type (or R-type) purinoceptors generally have the following characteristics: (1) show a potency sequence of Ado  $\geq$  AMP > ADP  $\geq$  ATP; (2) are most affected by structural changes in the ribose moiety of the stimulant; (3) prefer agonists in the *anti*-conformation; and (4) are antagonized by theophylline and other methylxanthines (52,55,56). Hence the only major difference between the  $P_1$ -type purinoceptor and  $P_1$ -like chemoreceptors of the lobster (and the grass shrimp) is that the chemoreceptor requires the presence of the 5'-phosphate group on the nucleoside. The 5'-phosphate group is apparently optional for activation of the  $P_1$ -type receptor of vertebrates.

In addition to the crustaceans described above, there are other animals possessing external chemoreceptors for certain nucleotides or nucleosides. The African armyworm, an insect larva, has a receptor for adenosine that may function in feeding behavior (57). Also, AMP is an attractant for larval mosquitos (58). Further, receptors on the lips of a fish, the puffer, respond preferentially to nucleotides, especially ADP, IMP, and UMP (59). Likewise, IMP and inosine are gustatory feeding stimulants in another fish, the turbot (60). Nucleotides have a different type of effect on external chemoreceptors found in bovine taste papillae. In this

case, the nucleotides GMP, IMP, and UMP increase the amount of glutamate bound to glutamate receptors (61).

## $P_2$ -like Purinergic Chemoreceptors

In addition to the  $P_1$ -like chemoreceptors described above, the spiny lobster has a second distinct population of purinergic chemoreceptors with response characteristics indicative of the  $P_2$ -type purinoceptors found in internal tissues of vertebrates (38). This second population is composed of ATP-sensitive cells with response properties easily distinguished from the AMP-sensitive ( $P_1$ -like) cells described earlier (Fig. 6). Responses given by the ATP-sensitive cells are very brief in duration (less than 0.5 sec), even at high concentrations, whereas the AMP-sensitive cells typically have response durations of 5 to 10 sec, or longer (Fig. 6). Also, the maximum discharges of the ATP-sensitive cells to ATP are of low magnitude (ca. 11 impulses/response, SEM  $\pm$  0.95) whereas the AMP-sensitive cells have high maximum discharges (ca. 104 impulses/response, SEM  $\pm$  24.5).

SAR studies of the ATP-sensitive cells were conducted with the nucleotides and other substances shown in Figure 7. In these cells, ATP is far more active than ADP, whereas AMP and Ado are virtually nonstimulatory. An activity sequence for other substances is as follows: ATP  $\geq$  deoxyATP (dATP) > GTP > CTP  $\geq$  XTP  $\geq$  ITP  $\gg$  8-bromo-ATP = ADP, with pyrophosphate and tripolyphosphate being virtually inactive. Hence, these cells possess a broad sensitivity to nucleotide triphosphates and tolerate changes in both the ribose and adenine moieties. The minor differences in activity seen with most nucleotide triphosphates contrasts dramatically with a ca. 20-fold decrease in activity seen when the diphosphate group of ADP is substituted for the triphosphate chain of ATP.

The ATP-sensitive cells are also strongly stimulated by three slowly degradable analogs of ATP, namely,  $\alpha,\beta$ -methylene ATP (AMPCPP),  $\beta,\gamma$ -methylene ATP (AMPPCP), and  $\beta,\gamma$ -imido ATP (AMPPNP), (see Fig. 7). The potency sequence is AMPPNP > ATP = AMPPCP > AMPCPP, with the analog AMPPNP being significantly more effective than ATP itself.

Collectively, the above results show that the ATP-sensitive cells in the antennules of the spiny lobster share the following properties with the  $P_2$ -type purinoceptors described by Burnstock (52). In the antennular receptors, ATP is far more effective than ADP, which in turn is more effective than either AMP or Ado. Secondly, the antennular receptors show a broad sensitivity to nucleotide triphosphates including those with modifications in both the adenine and ribose moieties. This broad sensitivity is consistent with that of  $P_2$ -type purinoceptors in brain (47) and in smooth muscle of various organs (62-64). Thirdly, both the antennular chemoreceptors and  $P_2$ -type purinoceptors are stimulated by the slowly degradable analogs of ATP, namely, AMPPNP, AMPPCP, and AMPCPP. Further, the analog AMPPNP is more effective than ATP itself on the



















Adenine AlterationsRibose Phosphate Alterations

FIGURE 5. Molecular formulae of AMP and analogs tested on AMP-sensitive chemosensory cells of the spiny lobster. Analogs with alterations in the adenine moiety (upper) and the ribose phosphate moiety (lower) were tested. The number beneath each substance represents the intensity of the physiological response to that substance (10  $\mu$ M) relative to the response to 10  $\mu$ M AMP. Data from Derby et al. (37).




FIGURE 6. Comparisons of response characteristics of ATP-sensitive cells and AMP-sensitive cells in the antennule of the spiny lobster. (A,B) Response profiles of the two cell types each tested at two concentrations. Note differences in time scale, plus differences in duration and magnitude of responses. (C,D) Dose-response curves for populations of each cell type. Data points are mean values ( $\pm$  SEM). From Carr et al. (38).

antennular receptors; this and other analogs are also more effective than ATP on certain  $P_2$  purinoceptors identified in internal tissues (65,66).

The role of ATP as an olfactory excitant in the spiny lobster is unknown. However, this substance does have a role as a gustatory stimulant that induces certain blood-sucking insects to gorge on liquid (67–69). Moreover, the behavioral responses evoked by ATP in these insects also appear to be mediated by a  $P_2$ -like purinoceptor (38).

## Inactivation and Uptake of Purinergic Substances

Further parallels between purinergic systems found in the lobster olfactory organ and in internal tissues of vertebrates are revealed by comparing the biochemical fate of excitatory nucleotides in both cases. In vertebrate tissues, the inactivation of extracellular nucleotides occurs by a two-step process: first, the nucleotides are dephosphorylated, and then the resultant Ado is internalized by an uptake system (70). Dephosphorylation is mediated by ectonucleotidases present on external membrane surfaces of neurons, glial cells, and other cell types (71–73). Ado is salvaged by an uptake system believed to involve facilitated diffusion (74–76).

In the spiny lobster, sensilla on the antennule also have an ectonucleotidase that inactivates the excitatory nucleotide, AMP, to yield the non-excitatory product, Ado, which is then internalized by an uptake system

within the sensilla (Trapido-Rosenthal et al., in preparation). Dephosphorylation is rapid; the rate of uptake of  $^3H$  from  $^3H$ -AMP (labeled in the purine ring) is indistinguishable from the rate of uptake of  $^3H$  from  $^3H$ -Ado (Fig. 8A). Evidence that AMP is indeed dephosphorylated prior to uptake is shown by the fact that the rate of uptake of  $^3H$  from ring-labeled AMP is about 10 times greater than that of  $^{32}P$  from  $^{32}P$ -AMP (Fig. 8B). Also, the 5'-nucleotidase inhibitor, AMP-C-P, inhibits the uptake of  $^3H$  from ring-labeled AMP but not from labeled Ado (Trapido-Rosenthal et al., in preparation).

## Taurinergic Chemoreceptors

### Background







Taurine (2-aminoethanesulfonic acid) is found at high concentrations in a wide variety of animal tissues (77) and has been the object of increased research effort during the past 15 years (78–82). This ubiquitous amino acid has been associated with neural development (83), heart function (84), retinal function (85), osmoregulation (86,87), opioid activity (88,89), and disease states such as epilepsy (90) and Friedreich's ataxia (91). Taurine has also been postulated to act as an inhibitory neurotransmitter and/or neuromodulator (92,93). Nevertheless, despite taurine's known or suspected involvement in numerous physiological processes, its functional significance remains largely speculative. Unlike other structurally related, bioactive amino acids (e.g., glycine and GABA), the role of taurine is poorly understood because of a paucity of selective agonists and antagonists for pharmacological studies. This lack of selective pharmacological probes is due in large measure to the absence of a model receptor system in vertebrates that is specifically sensitive to taurine (94). As described below, taurine-sensitive chemoreceptors of the lobster provide a unique model for exploring taurine reception in detail, and as such may prove invaluable in the identification of pharmacological agents for use in deciphering taurine's role in vertebrate systems.

### Lobster Taurinergic Chemoreceptors

Chemoreceptors narrowly tuned to taurine have been identified on the antennules of the spiny lobster, *P. argus* (33,36), and on the walking legs and antennules of the American lobster, *H. americanus* (26,35,95,96). In *P. argus*, average dose-response relationships for these olfactory cells show that the number of impulses produced, the maximum frequency of discharge, and the response duration all increase in an approximately linear fashion at concentrations between  $10^{-9}$  and  $10^{-4}$  M taurine (36). Although threshold differences exist between individual taurine-sensitive cells, single cells are consistent in their response to repeated stimulation. Threshold determinations have revealed responses to taurine at concentrations as low as  $10^{-12}$  M, the lowest dose examined (33).

Fuzessery et al. (36) investigated the stimulatory ca-

### Adenine Alterations



### Ribose Triphosphate Alterations











FIGURE 7. Molecular formulae of ATP and analogs tested on  $P_2$ -like chemoreceptors of the spiny lobster.






FIGURE 8. Biochemistry of purinergic substances in excised chemosensory sensilla of the spiny lobster. (A) Sensilla were incubated with 0.1  $\mu\text{M}$  adenine-labeled  $^3\text{H}$ -adenosine or  $^3\text{H}$ -AMP in the presence and absence of an excess (1000  $\mu\text{M}$ ) of unlabeled adenosine. The tritiated adenine moiety of AMP and adenosine were internalized at very similar rates. (B) Sensilla were incubated in a mixture containing both 0.1  $\mu\text{M}$   $^{32}\text{P}$ -AMP and 0.1  $\mu\text{M}$  adenine-labeled  $^3\text{H}$ -AMP in the presence and absence of excess unlabeled adenosine. The uptake of  $^3\text{H}$  but not  $^{32}\text{P}$  indicate that most AMP is dephosphorylated prior to uptake. From Trapido-Rosenthal et al., in preparation.

capacity of compounds structurally related to taurine (Fig. 9), each presented at a concentration of  $10^{-5}$  M. These data disclosed the following relationships [Note that numbers in brackets below refer to compound numbers in Fig. 9].

Compounds with one terminal basic group and one terminal acidic group separated by two carbon (C) atoms are the most active [1,2]. An exception occurs when a phosphonic acid group is substituted for the carboxyl group [3]. Compounds with the basic and acidic groups separated by greater than 2C atoms are less stimulatory with potency decreasing as a function of the distance of separation [e.g., 2>15>16>17]. Compounds with a 1C atom separation are not stimulatory [4,14].

Compounds lacking the terminal amino group [5,6,7,9,10] or with an alpha amine group in addition to the terminal amino group are virtually inactive [9,11,12]. Substituting a guanido group for the terminal amine results in loss of activity [20,21].

Neutral side chains markedly decrease the stimulatory capacity [15>18; 2>19].

Compounds with the carboxyl group blocked in a peptide bond are not stimulatory [22,23].

In addition, compounds structurally dissimilar to taurine [twelve  $\alpha$ -amino acids, three organic acids (citric, propionic and succinic) and the quaternary amine, glycine betaine] did not stimulate 65 other taurine-sensitive cells on which each compound was tested at  $10^{-5}$  M.

Collectively these findings indicate that the taurine receptor cells in *P. argus* have a very restricted specificity and exhibit interesting similarities in their response spectra to taurine-sensitive systems found in

internal tissues of vertebrates. For example, induced arrhythmias in dogs can be suppressed by taurine, but ethanesulfonic acid and other compounds which lack either the basic or acidic group are ineffective (97). In human blood platelets, uptake of taurine is competitively inhibited by  $\beta$ -alanine and hypotaurine, but not by the phosphonic acid analog (98). Likewise, hypotaurine and  $\beta$ -alanine compete with taurine for uptake in rat brain, while  $\alpha$ -amino acids do not (99,100). These results from vertebrate internal systems parallel those in the lobster olfactory system; i.e.,  $\beta$ -alanine and hypotaurine are the two most active analogs for the taurine-sensitive olfactory cells, whereas  $\alpha$ -amino acids, the phosphonic acid analog of taurine, and ethanesulfonic acid are all inactive.

Although not as extensively studied in *H. americanus*, taurine-sensitive chemoreceptors on the antennules (95,96) and walking legs (26,35) of this species appear to exhibit "narrow tuning" comparable to that found in *P. argus*. For example, in a sample of seven taurine-sensitive cells on the antennules of *H. americanus*, taurine was very effective, but a series of 14 other compounds (L-hydroxyproline, L-glutamate, L-glutamine, ammonium chloride, L-arginine, L-sucrose, ethanol, L-alanine, L-lysine, betaine, L-aspartate, glycine, L-leucine, and L-proline), each presented at a concentration of  $10^{-4}$  M, showed little or no stimulatory activity (96). Interestingly, when taurine was presented in mixture with all 14 of the above substances, the cells' responses were significantly suppressed. Further study of this phenomenon revealed that the suppression was concentration dependent; i.e., with diluted mixtures (each compound  $<10^{-6}$  M) cell responses were not sup-




FIGURE 9. Molecular formulae of compounds tested on taurine-sensitive chemoreceptors of spiny lobster. Indices of relative activity are as follows: (VA) very active; (A) active; (SA) slightly active; (I) virtually inactive. Numbers designating compounds are referred to in text. Modified from Fuzessery et al. (36).

pressed, and in fact tended to be slightly greater than with taurine presented alone (29).

The mechanism of the above mixture suppression is not understood. However, in *P. argus*, studies of suppression on a second type of taurine-sensitive chemosensory cell suggest that competition for common receptor sites may be involved (101). The taurine-sensitive cells exhibiting suppression are less sensitive to taurine than the cell population previously described for *P. argus*; these cells have thresholds close to  $10^{-5}$  M, and the dose-response functions have a lesser slope. In these cells, the response to taurine is inhibited when taurine is presented in binary mixture with certain amino acids. Glycine is one of the most effective inhibitors (suppressants), and an analysis of the nature of the inhibition has indicated a parallel shift to the right of the taurine dose-response function. These results are particularly intriguing in the light of taurine's suspected

interactions with glycine receptors in some vertebrate preparations (102–104).

## Glutamatergic Chemoreceptors

### Background

For many years the acidic amino acids, glutamate and aspartate, have been candidate excitatory neurotransmitters in both vertebrates and invertebrates (105–107). Studies with molluscs and coelenterates have indicated that receptors for these amino acids are present in the CNS and nerve-muscle preparations, respectively. Perhaps the most thoroughly investigated invertebrate group, however, has been the arthropods, where the excitatory potency of glutamate has been shown for neuromuscular junctions in both crustaceans and insects (105).

Although the excitatory effects of glutamate in the vertebrate CNS were described over 30 years ago (108), the characterization of receptors for this and other structurally related amino acids has progressed slowly. Current neuropharmacological data indicate that multiple receptor types exist for these excitatory compounds (107,109). At least three classes of synaptic receptors have been identified, each associated with a different ionophore. One class is activated preferentially by quisqualate and related compounds; L-glutamate and L-cysteate are candidate transmitters for these receptors. A second class is preferentially excited by *N*-methyl-D-aspartate (NMDA); L-aspartate may be a transmitter for this class, but is also an agonist for quisqualate receptors. The third receptor type is distinguished by its activation with kainic acid, the effects of which are unaffected by quisqualate or NMDA receptor antagonists. The endogenous transmitter(s) for the kainate class of receptors is unknown. In addition to these three receptor classes, there is emerging pharmacological evidence for other classes or subtypes, reflecting the unsettled state of understanding in this field (107,109).

### Lobster Glutamatergic Chemoreceptors

Glutamate sensitive chemoreceptors on both the antennules and walking legs have been described in a number of marine crustaceans (25,26,95,110–113). In single unit studies with antennule preparations, a range of cell specificities is apparent. For example, Derby and Ache (113) analyzed 70 receptor cells in *P. argus* for responses to eight substances including L-glutamate, taurine, betaine, L-alanine, glycine, AMP, L-proline, and hydroxy-L-proline, each presented at the relative concentrations at which they occur in a natural food stimulus mixture. Eighteen of the cells responded only to glutamate and betaine; 11 of these responded most strongly to glutamate, whereas the remaining seven cells exhibited approximately equal activation by glutamate and betaine. In *H. americanus*, only a single cell was encountered that responded exclusively to L-glutamate when 30 cells

were tested with a battery of 15 substances (96). A second glutamate-sensitive cell was equally stimulated by hydroxy-L-proline. Finally, it was reported that a single cell of nine examined in *H. americanus* responded only to L-glutamate when 29 D- and L-amino acids were tested (95).

In contrast to the antennular receptors described above, "narrowly tuned" glutamate chemoreceptors appear to comprise a major portion of the chemoreceptor population of the walking legs of the American lobster, *H. americanus* (26,35). In experiments examining the structure-activity relationships for activation of these cells, all of the glutamate analogs tested (Fig. 10) had a greatly reduced stimulatory capacity; i.e., responses were less than 8% of that to L-glutamate (26). These analogs included the following structural modifications [Note that numbers in brackets refer to compound numbers in Fig. 10]: (1) substituting phosphonic or sulfonic acid groups for the  $\gamma$ -carboxylic acid group [1,2]; (2)  $\alpha$ -carbon substitution [3]; (3) isomerization (D-glutamate); (4) deamination [4]; (5) cyclization [16]; (6) changing the carbon chain length [5,6]; (7) altering the  $\alpha$ -carboxylic acid group [7,8]; (8) alteration of the  $\gamma$ -carboxylic acid group [9,11,12,13]; (9) altering the carbon chain length as well as the  $\gamma$ -carboxylic acid group [14,15]; and (10) adding a second or third amino acid group [10,18]. Because cysteate [14], aspartate [5], and kainate [17] were all virtually inactive, these results suggest that the walking leg receptors represent a class that is distinctly different from the classes of internal "glutamate" receptors described earlier. Indeed this difference is apparent even within the same species. For example, aspartate potentiates the actions of L-glutamate at the neuromuscular synapse in *H. americanus*, presumably by modulating the affinity of the glutamate receptor (114); however, tests exploring similar potentiation with the glutamate chemoreceptors proved negative (26). Furthermore, unlike the glutamate chemoreceptor, the neuromuscular preparation in *H. americanus* is stimulated by kainate (115).

## Glycinergic Chemoreceptors

Glycine is known to act as an inhibitory neurotransmitter in vertebrates, and its role as such is best established in the spinal cord (116) where there is good evidence that Renshaw cells are glycinergic (117). Although glycine sensitive chemoreceptors have been found in crustaceans, their characterization has lagged behind other receptor types because of difficulties in locating single units. For example, even though glycine was one of the most stimulatory amino acids based on multiunit recordings in the walking legs of *H. americanus* (25), a search for single cells was unsuccessful (26). Also, within a sample group of 70 chemoreceptor cells on the antennules of *P. argus*, only two exhibited selective sensitivity to glycine (113). Recently, a new search protocol in our laboratory, using higher concentrations of glycine, has facilitated the identification of

glycine-sensitive cells, and these are currently under study.

Single chemoreceptive cells on crayfish (*Orconectes limosus*) legs have been characterized which are sensitive to a range of amino acids and related substances (118). These findings suggest that a single receptor site type, broadly sensitive to amino acids, mediates the responses of these cells. In this system glycine is moderately stimulatory with a  $K_m$  of  $1.5 \times 10^{-3}$  mole/L. This broad sensitivity to amino acids in a freshwater crustacean markedly contrasts with the situation for chemoreceptors of marine species, possibly indicating that the different ionic environments to which the chemoreceptors are exposed have given rise to the evolutionary divergence.

## Other Chemoreceptors

Antennular chemoreceptors of the California spiny lobster (*P. interruptus*) respond to  $10^{-9}$  to  $10^{-13}$  M concentrations of the steroid molting hormones,  $\alpha$ - and  $\beta$ -ecdysone (119). Chemoreceptors responding to  $\alpha$ -ecdysone do not respond to  $\beta$ -ecdysone, but additional details of receptor specificity are not known. The short latency period of these receptor cells indicates that the electrophysiological response is mediated by a membrane receptor rather than by a cytoplasmic or nuclear event such as characterizes the effects of many other steroids (119). Hence the antennular chemoreceptors may prove useful for comparative studies of steroid receptors located on cell surfaces.

Walking legs of crayfish (*Orconectes limosus* and *Austropotamobius torrentium*) have chemoreceptor cells that are sensitive to pyridine analogs but not to amino acids or amines (40,120). Tests of 79 pyridine analogs and related compounds on these cells revealed that the most effective substances were pyrazinecarboxamide > 3-acetylpyridine > nicotinamide > pyridine-3-aldoxime.  $K_m$  values for the above substances ranged from about  $10^{-6}$  to  $4 \times 10^{-4}$  M (28,121). 4-Acetylpyridine was identified as a competitive antagonist of the pyridine-sensitive cells. Although possible relationships between the pyridine-sensitive chemoreceptors of crayfish and internal receptor types have not been explored, the pyridine moiety is indeed represented in many substances known to have regulatory or metabolic effects in internal tissues (e.g., nicotinic acid, nicotinamide, NAD, NADP, and others) (122). Also, the pyridine-sensitive chemoreceptors are stimulated by 3- and 4-aminopyridines that are known to have excitatory effects on synaptic and axonal structures (123,124).

## Possible Evolutionary Implications of External Chemoreceptors Sensitive to Internal Neuroactive Agents

In multicellular aquatic organisms, the principal functions of external chemoreceptors and of internal recep-




FIGURE 10. Molecular formulae of compounds tested on glutamate-sensitive chemoreceptors of the American lobster. Numbers designating compounds are referred to in text. Formulae from Derby and Atema (26).

tors (e.g., synaptic receptors) are very similar in that both monitor particular chemicals in their surrounding aquatic environment. Both types of "chemo"-receptors have binding sites with an affinity for specific molecules, and both are coupled to a transduction mechanism capable of affecting membrane excitability. Indeed, Haldane (125) hypothesized that some internal receptor types may have evolved from external chemoreceptors of primitive aquatic animals. Support for this hypothesis is provided by the fact that all of the neuroactive sub-

stances listed in Table 1 are known to activate receptors existing both in internal tissues and on external chemosensory surfaces.

Further support for the hypothesis of a common origin for certain external and internal receptor types is indicated by the finding that receptors for the neurotransmitters acetylcholine, catecholamines, and GABA are variously distributed on the surface membranes of oocytes from frogs, mice, and humans (126,127). Such receptors are exposed to chemical signals present in an

Table 1. Neuroactive agents for which external chemoreceptors exist.

Substance	Organism with chemoreceptors for substance	Behavior influenced by substance	Reference
AMP	Shrimp, spiny lobster, mosquito larvae	Attraction of shrimp, mosquito larvae	(37, 53, 58)
Cyclic AMP	Slime mold	Attraction	(42, 129)
ATP	Mosquito, spiny lobster	Liquid ingestion by mosquito	(38, 68)
Acetylcholine ACH	Ciliated protozoan	Swimming velocity	(130)
$\gamma$ -Amino-butyric acid (GABA)	Abalone larvae	Settlement and metamorphosis	(131)
Dihydroxy-phenylalanine (DOPA)	Oyster larvae	Settlement and metamorphosis	(132)
Glutamate	Lobster, colonial algae	Induction of sexual stage in algae	(26, 133)
Glycine	Mud snail	Proboscis extension	(134)
Taurine	Spiny lobster	Attraction	(36)
$\alpha$ , $\beta$ -Ecdysone	Spiny lobster	Not known	(119)
Prostaglandin A <sub>2</sub> (PGA <sub>2</sub> )	Fish	Inhibition of feeding	(135)

external aquatic environment prior to the developmental events that distribute daughter cells into the internal layers of the embryo. In fact, the entire vertebrate nervous system with its many receptor types for transmitters and other neuroactive substances traces its embryological origin to an outer layer of cells, the ectoderm. The ectodermal cells interface with an external aquatic environment until the onset of gastrulation and the further developmental processes that create new internal aquatic interfaces between new neighboring cells, cell layers, and body fluids.

Certainly much can be learned by using the chemoreceptors of appropriate invertebrates as model systems for several receptor types found in internal tissues (37,128). Chemoreceptors of large crustaceans such as lobsters are particularly suitable for this purpose because the receptor cells are present on large, easily accessible, external appendages; i.e., the antennules and legs. More importantly, lobsters serve as excellent model systems because they are now known to have discrete chemoreceptors for at least six neuroactive agents, namely, AMP, ATP, glutamate, glycine, taurine and ecdysones (see Table 1). Moreover, techniques now exist to investigate the biochemical (Trapido-Rosenthal et al., in preparation) and biophysical (30) processes by which these agents activate chemoreceptor cells.

The fact that chemoreceptors are directly exposed to chemicals appearing in the external aquatic environment implies that the chemosensory system may be

particularly vulnerable to foreign chemicals or to other factors that modify the aquatic environment. Environmental variables that are known to influence certain chemosensory responses of aquatic animals include the following: temperature (136,137), pH (138), seasonality (139), feeding state (139), detergents (140), industrial wastes (141), drilling muds (142), oil (143), and certain heterocyclic compounds (144). Aside from recordings of the effects of temperature on pyridine- and amino acid-sensitive units (136,137), the effects of these environmental variables upon the discrete types of chemoreceptors described in this review are unknown.

We are grateful to Dr. Henry G. Trapido-Rosenthal for providing very recent data on the biochemistry of a purinergic chemosensory system. Ms. Marsha Lynn Milstead prepared the illustrations. This work was supported in part by NSF Grant Nos. BNS 84-11693 and BNS 85-11256.

## REFERENCES

1. Ache, B. W. Chemoreception and thermoreception. In: The Biology of Crustacea, Vol. 3 (H. L. Atwood and D. C. Sandemann, Eds.), Academic Press, New York, 1982, pp. 369-398.
2. Carr, W. E. S. Chemical stimulation of feeding behavior. In: Chemoreception in Fishes (T. J. Hara, Ed.), Elsevier Scientific Publishing Co., Amsterdam, 1982, pp. 259-275.
3. Mackie, A. M., and Grant, P. T. Interspecies and intraspecies chemoreception by marine invertebrates. In: Chemoreception in Marine Organisms (P. T. Grant and A. M. Mackie, Eds.), Academic Press, New York, 1974, pp. 105-141.
4. Sleeper, H. L., Paul, V. J., and Fenical, W. Alarm pheromones from the marine opisthobranch *Navanax inermis*. J. Chem. Ecol. 6: 57-70 (1980).
5. Atema, J., and Engstrom, D. G. Sex pheromone in the lobster, *Homarus americanus*. Nature 232: 261-263 (1971).
6. Gleeson, R. A. Pheromone communication in the reproductive behavior of the blue crab, *Callinectes sapidus*. Mar. Behav. Physiol. 7: 119-134 (1980).
7. Morse, D. E. Biochemical control of larval recruitment and fouling. In: Marine Biodeterioration: An Interdisciplinary Study (J. D. Costlow and R. C. Tipper, Eds.), Naval Institute Press, Annapolis, MD, 1984, pp. 134-140.
8. Larman, V. N. Protein extracts of some marine animals which promote barnacle settlement: possible relationship between a protein component of arthropod cuticle and actin. Comp. Biochem. Physiol. 77B: 73-81 (1984).
9. Hasler, A. D., Scholz, A. T., and Horrall, R. M. Olfactory imprinting and homing in salmon. Am. Scientist 66: 347-355 (1978).
10. Stabell, O. B. Homing and olfaction in salmonids: a critical review with special reference to the Atlantic salmon. Biol. Rev. 59: 333-388 (1984).
11. Todd, J. H., Atema, J., and Bardach, J. E. Chemical communication in social behavior of a fish, the yellow bullhead (*Ictalurus natalis*). Science 158: 672-673 (1967).
12. Liley, N. R. Chemical communication in fish. Can. J. Fish. Aquat. Sci. 39: 22-35 (1982).
13. Daloz, D., Braekman, J. C., and Tursch, B. Chemical communication in the marine environment. In: Animals and Environmental Fitness. Physiological and Biochemical Aspects of Adaptation and Ecology, Vol. 1 (R. Gilles, Ed.), Pergamon Press, Oxford, 1980, pp. 243-261.
14. Atema, J. Chemoreception in the sea: adaptations of chemoreceptors and behavior to aquatic stimulus conditions. Soc. Exp. Biol. Sympos. 39: 387-423 (1985).
15. Carr, W. E. S. The molecular nature of chemical stimuli in the aquatic environment. In: Sensory Biology of Aquatic Animals (J. Atema, R. R. Fay, A. N. Popper, and W. N. Tavolga, Eds.), Springer-Verlag, New York, 1987.

16. Ache, B. W., and Derby, C. D. Functional organization of olfaction in crustaceans. *Trends NeuroSci.* 8: 356-360 (1985).
17. Laverack, M. S., and Ardill, D. J. The innervation of the aesthetasc hairs of *Panulirus argus*. *Quart. J. Micr. Sci.* 106: 45-60 (1965).
18. Sandeman, D. C., and Luff, S. E. The structural organization of glomerular neuropile in the olfactory and accessory lobes of an Australian freshwater crayfish, *Cherax destructor*. *Z. Zellforsch. Mikrosk. Anat.* 142: 37-61 (1973).
19. Sandeman, D. C., and Denberg, J. L. The central projections of chemoreceptor axons in the crayfish revealed by axoplasmic transport. *Brain Res.* 115: 492-496 (1976).
20. Derby, C. D. Structure and function of cuticular sensilla of the lobster *Homarus americanus*. *J. Crust. Biol.* 2: 1-21 (1982).
21. Altner, H., and Prillinger, L. Ultrastructure of invertebrate chemo-, thermo-, and hygroreceptors and its functional significance. *Int. Rev. Cytol.* 67: 69-139 (1980).
22. Hamilton, K. A. A topographical and typographical comparison of the rodlike setae of ambulatory dactylopodites in decapod crustaceans. *J. Morphol.* 176: 351-364 (1983).
23. Hamilton, K. A., Linberg, K. A., and Case, J. F. Structure of dactyl sensilla in the kelp crab, *Pugettia producta*. *J. Morphol.* 185: 349-366 (1985).
24. Atema, J. Functional separation of smell and taste in fish and crustacea. In: *Olfaction and Taste*, Vol. VI (J. LeMagnen and P. MacLeod, Eds.), IRL Press, London, 1977, pp. 165-174.
25. Derby, C. D., and Atema, J. Chemosensitivity of walking legs of the lobster *Homarus americanus*: neurophysiological response spectrum thresholds. *J. Exptl. Biol.* 98: 303-315 (1982).
26. Derby, C. D., and Atema, J. Narrow-spectrum chemoreceptor cells in the walking legs of the lobster *Homarus americanus*: taste specialists. *J. Comp. Physiol.* 146: 181-189 (1982).
27. Hatt, H. Structural requirements of amino acids and related compounds for stimulation of receptors in crayfish walking leg. *J. Comp. Physiol.* 155: 219-231 (1984).
28. Hatt, H., and Schmiedel-Jakob, I. Electrophysiological studies of pyridine-sensitive units on the crayfish walking leg. I. Characteristics of stimulatory molecules. *J. Comp. Physiol.* 154: 855-863 (1984).
29. Johnson, B. R., Borroni, P. F., and Atema, J. Mixture effects in primary olfactory and gustatory receptor cells from the lobster. *Chem. Senses* 10: 367-373 (1985).
30. Anderson, P. A. V., and Ache, B. W. Voltage- and current-clamp recordings of the receptor potential in olfactory receptor cells in situ. *Brain Res.* 338: 273-280 (1985).
31. Schmitt, B. C., and Ache, B. W. Olfaction: responses of a decapod crustacean are enhanced by flicking. *Science* 205: 204-206 (1979).
32. Seelinger, G. Response characteristics and specificity of chemoreceptors in *Hemilepistus reaumuri* (Crustacea, Isopoda). *J. Comp. Physiol.* 152: 219-229 (1983).
33. Thompson, H., and Ache, B. W. Threshold determination for olfactory receptors of the spiny lobster. *Mar. Behav. Physiol.* 7: 249-260 (1980).
34. Derby, C. D., and Atema, J. Chemoreceptor cells in aquatic invertebrates: peripheral mechanisms of chemical signal processing in decapod crustaceans. In: *Sensory Biology of Aquatic Animals* (J. Atema, R. R. Fay, A. N. Popper, and W. N. Tavolga, Eds.), Springer-Verlag, New York, 1987.
35. Johnson, B. R., Voigt, R., Borroni, P. F., and Atema, J. Response properties of lobster chemoreceptors: tuning of primary taste neurons in walking legs. *J. Comp. Physiol.* 155: 593-604 (1984).
36. Fuzessery, Z. M., Carr, W. E. S., and Ache, B. W. Antennular chemosensitivity in the spiny lobster, *Panulirus argus*: studies of taurine sensitive receptors. *Biol. Bull.* 154: 226-240 (1978).
37. Derby, C. D., Carr, W. E. S., and Ache, B. W. Purinergic olfactory receptors of lobsters are similar to internal purinergic receptors of vertebrates. *J. Comp. Physiol.* 155: 341-349 (1984).
38. Carr, W. E. S., Gleeson, R. A., Ache, B. W., and Milstead, M. L. Olfactory receptors of the spiny lobster: ATP-sensitive cells with similarities to P2-type purinoceptors of vertebrates. *J. Comp. Physiol.* 158: 331-338 (1986).
39. Schneider, D., Lacher, V., and Kaissling, K. E. Die Reaktionsweise und das Reaktionsspektrum von Riechzellen bei *Antheraea pernyi* (Lepidoptera, Saturniidae). *Z. Vergl. Physiol.* 48: 632-662 (1964).
40. Bauer, U., and Hatt, H. Demonstration of three different types of chemosensitive units in the crayfish claw using a computerized evaluation. *Neurosci. Lett.* 17: 209-214 (1980).
41. Sass, H. Olfactory receptors on the antenna of *Periplaneta*: response constellations that encode food odors. *J. Comp. Physiol.* 128: 227-233 (1978).
42. Mato, J. M., Jastorff, B., Morr, M., and Konijn, T. M. A model for cyclic AMP-chemoreceptor interaction in *Dictyostelium discoideum*. *Biochem. Biophys. Acta* 544: 309-314 (1978).
43. Greengard, P. C. Intracellular signals in the brain. The Harvey Lectures, Series 75, Academic Press, New York, 1981, pp. 277-331.
44. Berridge, M. J. The molecular basis of communication within the cell. *Sci. American* 253: 142-152 (1985).
45. Burnstock, G., and Brown, C. M. An introduction to purinergic receptors. In: *Purinergic Receptors* (G. Burnstock, Ed.), Chapman and Hall, London, 1981, pp. 1-45.
46. Stone, T. W. Physiological roles for adenosine and adenosine 5'-triphosphate in the nervous system. *Neuroscience* 6: 523-555 (1981).
47. Phillis, J. W., and Wu, P. H. The role of adenosine and its nucleotides in central synaptic transmission. *Progr. Neurobiol.* 16: 187-239 (1981).
48. Daly, J. W. Role of ATP and adenosine receptors in physiologic processes: summary and prospectus. In: *Physiology and Pharmacology of Adenosine Derivatives* (J. W. Daly, Y. Kuroda, J. W. Phillis, H. Shimizu, and M. Ui, Eds.), Raven Press, New York, 1983, pp. 275-290.
49. Su, C. Purinergic neurotransmission and neuromodulation. *Ann. Rev. Pharmacol. Toxicol.* 23: 397-411 (1983).
50. Burnstock, G., and Buckley, N. J. The classification of receptors for adenosine and adenine nucleotides. In: *Methods in Pharmacology*, Vol. 6 (D. M. Paton, Ed.), Plenum Press, 1985, pp. 193-212.
51. Snyder, S. H. Adenosine as a neuromodulator. *Ann. Rev. Neurosci.* 8: 103-124 (1985).
52. Burnstock, G. A basis for distinguishing two types of purinergic receptors. In: *Cell Membrane Receptors for Drugs and Hormones: A Multidisciplinary Approach* (R. W. Straub and L. Bolis, Eds.), Raven Press, New York, 1978, pp. 107-118.
53. Carr, W. E. S., and Thompson, H. Adenosine 5'-monophosphate, an internal regulatory agent, is a potent chemoattractant for a marine shrimp. *J. Comp. Physiol.* 153: 47-53 (1983).
54. Lee, C. H., Evans, F. E., and Sarma, R. H. Interrelation between glycosidic torsion, sugar pucker, and backbone conformation in 5'- $\beta$ -nucleotides. *J. Biol. Chem.* 250: 1290-1296 (1975).
55. Londos, C., and Wolff, J. Two distinct adenosine-sensitive sites on adenylate cyclase. *Proc. Natl. Acad. Sci. (U.S.)* 74: 5482-5486 (1977).
56. Londos, C., Cooper, D. M. F., and Wolff, J. Subclasses of external adenosine receptors. *Proc. Natl. Acad. Sci. (U.S.)* 77: 2551-2554 (1980).
57. Ma, W. C. Electrophysiological evidence for chemosensitivity to adenosine, adenine and sugars in *Spodoptera exempta* and related species. *Experientia* 33: 356-358 (1977).
58. Barber, J. T., Ellgard, E. G., and Herskowitz, K. The attraction of larvae of *Culex pipiens quinquefasciatus* Say to ribonucleic acids and nucleotides. *J. Insect Physiol.* 28: 585-588 (1982).
59. Kiyohara, S., Hidaka, I., and Tamura, T. Gustatory response in the puffer. II. Single fiber analyses. *Bull. Japan. Soc. Sci. Fish.* 41: 383-391 (1975).
60. Mackie, A. M., and Adron, J. W. Identification of inosine and inosine 5'-monophosphate as the gustatory feeding stimulants for the turbot, *Scophthalmus maximus*. *Comp. Biochem. Physiol.* 60A: 79-83 (1978).
61. Torii, K., and Cagan, R. H. Biochemical studies of taste sensation. IX. Enhancement of L-[ $^3$ H] glutamate binding to bovine taste papillae by 5'-ribonucleotides. *Biochim. Biophys. Acta* 627: 313-323 (1980).

62. Maguire, M. H., and Satchell, D. G. Specificity of adenine nucleotide receptor sites: inhibition of the guinea pig taenia coli by adenine nucleotide analogs. In: *Physiological and Regulatory Functions of Adenosine and Adenine Nucleotides* (H. P. Baer and G. I. Drummond, Eds.), Raven Press, New York, 1979, pp. 33-43.
63. Brown, C. M., and Burnstock, G. Evidence in support of the P1/P2 purinoceptor hypothesis in the guinea-pig taenia coli. *Brit. J. Pharmacol.* 73: 617-624 (1981).
64. Lukacszo, P., and Krell, R. D. Responses of guinea-pig urinary bladder to purine and pyrimidine nucleotides. *Eur. J. Pharmacol.* 80: 401-406 (1982).
65. Kennedy, C., and Burnstock, G. ATP produces vasodilation via P1 purinoceptors and vasoconstriction via P2 purinoceptors in the isolated rabbit central ear artery. *Blood Vessels* 22: 145-155.
66. Kennedy, C., Delbro, D., and Burnstock, G. P2-purinoceptors mediate both vasodilation (via the endothelium) and vasoconstriction of the isolated rat femoral artery. *Eur. J. Pharmacol.* 107: 161-168 (1985).
67. Smith, J. J. B., and Friend, W. G. Further studies on potencies of nucleotides as gorging stimuli during feeding in *Rhodnius prolixus*. *J. Insect Physiol.* 22: 607-611 (1976).
68. Friend, M. G., and Smith, J. J. B. Factors affecting feeding by bloodsucking insects. *Ann. Rev. Entomol.* 22: 309-331 (1977).
69. Friend, M. G., and Smith, J. J. B. ATP analogs and other phosphate compounds as gorging stimulants for *Rhodnius prolixus*. *J. Insect Physiol.* 28: 371-376 (1982).
70. Burnstock, G. Comparative studies of purinergic nerves. *J. Exptl. Zool.* 194: 103-134 (1975).
71. Trams, E. G. Evidence for ATP action on the cell surface. *Nature* 252: 480-482 (1974).
72. Kreutzberg, G. W., Barron, K. D., and Schubert, P. Cytochemical localization of 5'-nucleotidase in glial plasma membranes. *Brain Res.* 158: 247-257 (1978).
73. Pearson, J. D., Carleton, J. S., and Gordon, J. L. Metabolism of adenine nucleotides by ectoenzymes of vascular endothelial and smooth-muscle cells in culture. *Biochem. J.* 190: 421-429 (1980).
74. Hertz, L. Kinetics of adenosine uptake into astrocytes. *J. Neurochem.* 31: 55-62 (1978).
75. Bender, A. S., Wu, P. H., and Phillis, J. W. The rapid uptake and release of [<sup>3</sup>H]adenosine by rat cerebral synaptosomes. *J. Neurochem.* 36: 651-660 (1981).
76. Kreutzberg, G. W., Reddington, M., Lee, K. S., and Schubert, P. Adenosine: transport, function and interaction with receptors in the CNS. *J. Neural Transmis. Suppl.* 18: 113-119 (1983).
77. Jacobsen, J. G., and Smith, L. H. Biochemistry and physiology of taurine and taurine derivatives. *Physiol. Rev.* 48: 424-511 (1968).
78. Huxtable, R. J., and Barbeau, A., Eds. *Taurine*. Raven Press, New York, 1976, p. 398.
79. Barbeau, A., and Huxtable, R. J., Eds. *Taurine and Neurological Disorders*. Raven Press, New York, 1978, p. 468.
80. Schaffer, S. W., Baskin, S. L., and Koches, J. J., Eds. *Monographs of the Physiological Society of Philadelphia*, Vol. 7. S. P. Medical and Scientific Books, New York, 1981, p. 446.
81. Huxtable, R. J., and Pasantes-Morales, H., Eds. *Taurine in Nutrition and Neurology*. Plenum Press, New York, 1982, p. 551.
82. Kuriyama, K., Huxtable, R. J., and Iwata, H., Eds. *Sulfur Amino Acids: Biochemical and Clinical Aspects*. Alan R. Liss Inc., New York, 1983, p. 485.
83. Gaull, G. E., and Rassin, D. K. Taurine and brain development: human and animal correlates. In: *Neural Growth and Differentiation* (E. Meisami and M. A. B. Brazier, Eds.), Raven Press, New York, 1979, pp. 461-477.
84. Grosso, D. A., and Bressler, R. Taurine and cardiac physiology. *Biochem. Pharmacol.* 25: 2227-2232 (1976).
85. Pasantes-Morales, H., Kleithi, J., Urban, P. F., and Mandel, P. The physiological role of taurine in retina: uptake and effect on electroretinogram (ERG). *Physiol. Chem. Phys.* 4: 339-348 (1972).
86. Simpson, J., Allen, K., and Awapara, J. Free amino acids in some aquatic invertebrates. *Biol. Bull.* 117: 371-381 (1959).
87. Thurston, J. H., Hauhart, R. E., and Naccarato, E. F. Taurine: possible role in osmotic regulation of mammalian heart. *Science* 214: 1373-1374 (1981).
88. Izumi, K., Munekata, E., Yamamoto, H., Nakanishi, T., and Barbeau, A. Effects of taurine and  $\gamma$ -aminobutyric acid on akinesia and analgesia induced by D-Ala<sup>2</sup>-Met-enkephalinamide in rats. *Peptides* 1: 139-146 (1980).
89. Yamamoto, H., McCain, H. W., Izumi, K., Misawa, S., and Way, E. L. Effects of amino acids, especially taurine and  $\gamma$ -aminobutyric acid (GABA) on analgesia and calcium depletion induced by morphine in mice. *Eur. J. Pharmacol.* 71: 177-184 (1981).
90. van Gelder, N. M., Sherwin, A. L., Sacks, C., and Andermann, F. Biochemical observations following administration of taurine to patients with epilepsy. *Brain Res.* 94: 297-306 (1975).
91. Barbeau, A., Melancon, S., Huxtable, R. J., and Lemieux, B. Taurine and Friedreich's ataxia; an update. In: *Taurine in Nutrition and Neurology* (Adv. Exp. Med., Vol. 139, R. Huxtable and H. Pasantes-Morales, Eds.), Plenum Press, New York, 1982, pp. 389-399.
92. Davison, A. N., and Kaczmarek, L. K. Taurine: a possible neurotransmitter. *Nature* 234: 107-108 (1971).
93. Mandel, P., Pasantes-Morales, H., and Urban, P. F. Taurine: a putative transmitter in retina. In: *Transmitters in the Visual Process* (S. L. Bonting, Ed.), Pergamon Press, New York, 1976, pp. 89-105.
94. Barbeau, A. Does taurine have clinical significance? In: *Taurine in Nutrition and Neurology* (Adv. Exp. Med., Vol. 139, R. Huxtable and H. Pasantes-Morales, Eds.), Plenum Press, New York, 1982, pp. 513-532.
95. Shephard, P. Chemoreception in the antennule of the lobster, *Homarus americanus*. *Mar. Behav. Physiol.* 2: 261-273 (1974).
96. Johnson, B. R., and Atema, J. Narrow-spectrum chemoreceptor cells in the antennules of the American lobster. *Neurosci. Lett.* 41: 145-150 (1983).
97. Welty, J. D., Read, W. O., and Byington, K. H. Comparison of amino-sulfonic acids as antiarrhythmic agents in dogs. In: *Taurine* (R. Huxtable and A. Barbeau, Eds.), Raven Press, New York, 1976, pp. 169-173.
98. Gant, Z. N., and Nauss, C. B. Uptake of taurine by human blood platelets: a possible model of the brain. In: *Taurine* (R. Huxtable and A. Barbeau, Eds.), Raven Press, New York, 1976, pp. 99-121.
99. Kaczmarek, L. K., and Davison, A. N. Uptake and release of taurine from rat brain slices. *J. Neurochem.* 19: 2355-2362 (1972).
100. Lähdesmäki, P., and Oja, S. S. On the mechanism of taurine transport at the brain cell membranes. *J. Neurochem.* 20: 1411-1417 (1973).
101. Gleeson, R. A., and Ache, B. W. Amino acid suppression of taurine-sensitive chemosensory neurons. *Brain Res.* 335: 99-107 (1985).
102. Curtis, D. R., Holsi, L., and Johnston, G. A. R. A pharmacological study of the depression of spinal neurons by glycine and related amino acids. *Exptl. Brain Res.* 6: 1-18 (1968).
103. Haas, H. L., and Hosli, L. The depression of brainstem neurons by taurine and its interaction with strychnine and bicuculline. *Brain Res.* 52: 399-402 (1979).
104. Kurachi, M., Yoshihara, K., and Aihara, H. Effect of taurine on depolarizations induced by L-glutamate and other excitatory amino acids in the isolated spinal cord of the frog. *Japan. J. Pharmacol.* 33: 1247-1254 (1983).
105. Usherwood, P. N. R. Amino acids as neurotransmitters. *Adv. Comp. Physiol. Biochem.* 7: 227-309 (1978).
106. Di Chiara, G., and Gessa, G. L., Eds. *Glutamate as a Neurotransmitter*. Raven Press, New York, 1981, p. 445.
107. McLennan, H. Receptors for the excitatory amino acids in the mammalian central nervous system. *Progr. Neurobiol.* 20: 251-271 (1983).
108. Hayashi, T. A physiological study of epileptic seizures following cortical stimulation in animals and its application to human clinics. *Japan. J. Physiol.* 3: 46-64 (1952).

109. Fagg, G. E. L-Glutamate, excitatory amino acid receptors and brain function. *Trends NeuroSci.* 8: 207-210 (1985).
110. Case, J., and Gwilliam, G. F. Amino acid sensitivity of the dactyl chemoreceptors of *Carcinides maenas*. *Biol. Bull.* 121: 449-455 (1961).
111. Johnson, B. R., and Ache, B. W. Antennular chemosensitivity in the spiny lobster, *Panulirus argus*: amino acids as feeding stimuli. *Mar. Behav. Physiol.* 5: 145-157 (1978).
112. Nolen, T. G. Chemoreception in the rock crab, *Cancer antennarius*: specificity of the dactyl glutamate chemoreceptors. Master of Arts Thesis, University of California, Santa Barbara, 1978, p. 147.
113. Derby, C. D., and Ache, B. W. Quality coding of a complex odorant in an invertebrate. *J. Neurophysiol.* 51: 906-924 (1984).
114. Freeman, A. R., Shank, R. P., Kephart, J., Dekin, M., and Wang, M. Modulation of transmission at a glutamate synapse. *J. Physiol. (Paris)* 75: 605-610 (1979).
115. Shank, R. P., and Freeman, A. R. Agonistic and antagonistic activity of glutamate analogs on neuromuscular excitation in the walking limbs of lobsters. *J. Neurobiol.* 7: 23-36 (1976).
116. Aprison, M. H., and Nadi, N. S. Glycine: inhibition from the sacrum to the medulla. In: *Amino Acids as Chemical Transmitters* (F. Fonnum, Ed.), Plenum Press, New York, 1978, pp. 531-570.
117. Lodge, D., Curtis, D. R., and Brand, S. J. A pharmacological study of the inhibition of ventral group I  $\alpha$ -excited spinal interneurons. *Exptl. Brain Res.* 29: 97-105 (1977).
118. Bauer, U., Dudel, J., and Hatt, H. Characteristics of single chemoreceptive units sensitive to amino acids and related substances in the crayfish leg. *J. Comp. Physiol.* 144: 67-74 (1981).
119. Spencer, M., and Case, J. F. Exogenous ecdysteroids elicit low threshold sensory responses in spiny lobsters. *J. Exptl. Zool.* 229: 163-166 (1984).
120. Hatt, H., and Bauer, U. Electrophysiological properties of pyridine receptors in the crayfish walking leg. *J. Comp. Physiol.* 148: 221-224 (1982).
121. Hatt, H., and Schmiedel-Jakob, I. Specific antagonists at the pyridine receptor: evidence from electrophysiological studies with acetylpyridines. *Chem. Senses* 10: 317-323 (1985).
122. Marcus, R., and Coulson, A. M. Water-soluble vitamins. In: *The Pharmacological Basis of Therapeutics*, 7th Ed. (A. G. Gilman, L. S. Goodman, T. W. Rall, and F. Murad, Eds.), MacMillan Publishing Co., New York, 1985, pp. 1551-1572.
123. Thesleff, S. Aminopyridines and synaptic transmission. *Neuroscience* 5: 1413-1419 (1980).
124. Hermann, A., and Gorman, A. L. F. Effects of 4-aminopyridine on potassium currents in a molluscan neuron. *J. Gen. Physiol.* 78: 63-86 (1981).
125. Haldane, J. B. S. La signalisation animale. *Anee Biol.* 58: 89-98 (1954).
126. Kusano, K., Miledi, R., and Stinnakre, J. Cholinergic and catecholaminergic receptors in the *Xenopus* oocyte membrane. *J. Physiol. (London)* 328: 143-170 (1982).
127. Dolci, S., Eusebi, F., and Siracusa, G.  $\gamma$ -Amino butyric-*n*-acid sensitivity of mouse and human oocytes. *Dev. Biol.* 109: 242-246 (1985).
128. Lenhoff, H. M., and Heagy, W. Aquatic invertebrates: model systems for study of receptor activation and evolution of receptor proteins. *Ann. Rev. Pharmacol. Toxicol.* 17: 243-258 (1977).
129. Van Haastert, P. J. M., Jastorff, B., Pinas, J. E., and Konijn, T. M. Analogs of cyclic AMP as chemoattractants and inhibitors of *Dictyostelium* chemotaxis. *J. Bacteriol.* 149: 99-105 (1982).
130. Brown, I. D., and Kerkut, G. A. A study of the chemokinetic effects of various pharmacological agents upon *Tetrahymena vorax*. *Comp. Biochem. Physiol.* 69C: 275-280 (1981).
131. Morse, D. E., Hooker, N., Duncan, H., and Jensen, L.  $\gamma$ -Aminobutyric acid, a neurotransmitter, induces planktonic abalone larvae to settle and begin metamorphosis. *Science* 204: 407-410 (1979).
132. Coon, S. L., Bonar, D. B., and Weiner, R. M. Induction of settlement and metamorphosis of the Pacific oyster, *Crassostrea gigas* (Thunberg), by L-DOPA and catecholamines. *J. Exptl. Mar. Biol. Ecol.* 94: 211-221 (1985).
133. Maier, I., and Muller, D. G. Sexual pheromones in algae. *Biol. Bull.* 170: 145-175 (1986).
134. Carr, W. E. S. Chemoreception in the mud snail, *Nassarius obsoletus*. II. Identification of stimulatory substances. *Biol. Bull.* 133: 106-127 (1967).
135. Gerhart, D. J. Prostaglandin A2: an agent of chemical defense in the Caribbean gorgonian *Plexaura homomalla*. *Mar. Ecol. Progr. Ser.* 19: 181-187 (1984).
136. Hatt, H. Temperature dependence of the response of the pyridine-sensitive units in the crayfish walking leg. *J. Comp. Physiol.* 152: 395-403 (1983).
137. Hatt, H. Effect of temperature on the activity of the amino acid receptors in the crayfish walking leg. *J. Comp. Physiol.* 152: 405-409 (1983).
138. Hara, T. J. Effects of pH on the olfactory responses to amino acids in rainbow trout, *Salmo gairdneri*. *Comp. Biochem. Physiol.* 54A: 37-39 (1976).
139. Carr, W. E. S., Netherton, J. C. III, and Milstead, M. L. Chemoattractants of the shrimp, *Palaemonetes pugio*: variability in responsiveness and the stimulatory capacity of mixtures containing amino acids, quaternary ammonium compounds, purines and other substances. *Comp. Biochem. Physiol.* 77A: 469-474 (1984).
140. Bardach, J. E., Fujiya, M., and Holl, A. Detergents: effects on the chemical senses of the fish *Ictalurus natalis* (JeSeur). *Science* 148: 1605-1607 (1965).
141. Johnstone, A. D. F., Whitford, D., and Hawkins, A. D. The effects of an industrial waste (cryolite recovery sludge) upon the Atlantic salmon, *Salmo salar* (L.). *Water Res.* 16: 1529-1535 (1982).
142. Derby, C. D., and Atema, J. Influence of drilling muds on the primary chemosensory neurons in the walking legs of the lobster, *Homarus americanus*. *Can. J. Fish. Aquat. Sci.* 38: 268-274 (1981).
143. Atema, J., Karnofsky, E. B., and Oleszko-Szuts, S. Lobster behavior and chemoreception: sublethal effects of No. 2 fuel oil. In: *Advances in Marine Environment Research* (F. S. Jacoff, Ed.), Environmental Research Laboratory, Office of Research and Development, U.S. Environmental Protection Agency, Narragansett, Rhode Island, 1979, pp. 122-134.
144. Hara, T. J. Is morpholine an effective olfactory stimulant in fish? *J. Fish. Res. Board Can.* 31: 1547-1550 (1974).